Check List the journal of biodiversity data

NOTES ON GEOGRAPHIC DISTRIBUTION

Check List 17 (1): 151–157
https://doi.org/10.15560/17.1.151

PENSOFT

Current distribution of the invasive mollusk *Corbicula fluminea* (O.F. Müller, 1774) (Bivalvia, Cyrenidae) in Brazil, including a new record from the state of Piauí

Manuella Feitosa Leal[1*], Luiz Ricardo Lopes de Simone[2], Ana Carolina Figueiredo Lacerda[1], Edson Lourenço da Silva[3], Tamaris Gimenez Pinheiro[4]

1 Programa de Pós-Graduação em Biologia (Zoologia), Departamento de Sistemática e Ecologia, Centro de Ciências exatas e da Natureza, Universidade Federal da Paraíba, João Pessoa, PB, Brazil • MFL: manuellafeitosa94@gmail.com https://orcid.org/0000-0002-5139-8616 • ACFL: acflacerda@dse.ufpb.br https://orcid.org/0000-0001-9014-9105

2 Universidade de São Paulo Museu de Zoologia, São Paulo, SP, Brazil • lrsimone@usp.br https://orcid.org/0000-0002-1397-9823

3 Instituto Federal do Piauí, Picos, Piauí, Brazil • ed.loren@ifpi.edu.br https://orcid.org/0000-0002-4442-5137

4 Curso de Licenciatura em Educação do Campo/Ciências da Natureza, Universidade Federal do Piauí, Picos, Piauí, Brazil • tamarisgimenez@gmail.com https://orcid.org/0000-0001-7246-2691

* Corresponding author

Abstract

Corbicula fluminea is an invasive clam originally from Asia that threatens the biodiversity of freshwater communities around the world. It has greatly expanded in Brazil since its first record. We report here the occurrence of *C. fluminea* from Piauí state, Parnaíba river basin and, provide an assessment of its current distribution in Brazil. The compiled data include 382 records in 22 states and the Federal District, encompassing a variety of ecosystems. These results contribute to a discussion about the invasion and expansion process of *C. fluminea* in Brazil over time.

Keywords

Bioinvasion, freshwater bivalve, semi-arid region

Academic editor: Bárbara Romera | Received 7 July 2020 | Accepted 16 January 2021 | Published 1 February 2021

Citation: Leal MF, Simone LRL, Lacerda ACF, Silva EL, Pinheiro TG (2021) Current distribution of the invasive mollusk *Corbicula fluminea* (O.F. Müller, 1774) (Bivalvia, Cyrenidae) in Brazil, including a new record from the state of Piauí. Check List 17 (1): 151–157. https://doi.org/10.15560/17.1.151

Introduction

Invasive species are characterized by their ability to cause ecological and economic impacts on ecosystems outside of their native range (Lockwood et al. 2007). Generally, such invaders exhibit opportunistic behavior, allocating a large amount of energy for reproduction (r-strategists), which can occur several times a year (Crespo et al. 2015). In addition to the inherent attributes of invasive species, successful invasions depend on the mechanism of introduction and features of the invaded environment (Sousa et al. 2008; Simberloff 2009; Crespo et al. 2015).

Biological invasions are among the main threats to the biodiversity of freshwater ecosystems, and some freshwa-

©The authors. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ter invasive mussels are among the most harmful organisms. This is especially true for *Corbicula fluminea* (O.F. Müller, 1774) (Sousa et al. 2008), a species originally from Southeast Asia and currently found on all continents (Poleze and Callil 2015; Gama et al. 2017). The first occurrence of *C. fluminea* in Brazil was recorded in the Rio Grande do Sul state in 1979 (Veitenheimer-Mendes 1981) and, since then, it has been recorded in all major hydrographic basins of the country (Santana et al. 2013). In the Northeast Region, it was recorded in the states of Bahia, Maranhão, Paraíba, Pernambuco, Rio Grande do Norte, and Sergipe (Santana et al. 2013; Almeida et al. 2015). Besides *C. fluminea*, three additional congeners have invaded Brazil: *C. fluminalis* (O.F. Müller, 1774), *C. largillierti* (Philippi, 1844), and *Corbicula* sp., and may have been misidentified as *C. fluminea*. Accounts on the identification and taxonomy can be found in Santos et al. (2012).

Corbicula fluminea presents high invasive potential and adaptive success in new environments due to rapid growth, early sexual maturation, short life cycle, high tolerance to different abiotic conditions, and dispersal ability. *Corbicula fluminea* is commonly dispersed through birds (but also through other animals), water streams, or by human action (such as ballast water, sand transportation, and fishkeeping) (Poleze and Callil 2015; Gama et al. 2016; Gomes et al. 2016). These features make *C. fluminea* one of the main agents responsible for economic losses in hydroelectric and water treatment plants. High population densities in these locations commonly cause duct clogging and increased corrosion rate in pipes (Ilarri and Sousa 2012; Gama et al. 2017).

Ecologically, the invasion by *C. fluminea* changes the structure and the function of ecosystems and biotic interactions by forming highly dense populations (Ilarri and Sousa 2012). Such changes are mainly caused by the mussel's high suspended solid filtration capacity, which influences light penetration and enacts subsequent changes in sediment composition, substrate-water column oxygenation dynamics, and nutrient cycling (Ilarri and Sousa 2012; Santana et al. 2013; Gomes et al. 2016). These environmental alterations affect the development of other organisms such as algae, freshwater sponges, crustaceans, insects, and gastropods (Sousa et al. 2008). *Corbicula fluminea* can affect native freshwater mussels through displacement or reduction of habitats available to juveniles. Competition for food, and ingestion of sperm, larvae (glochids), and juvenile individuals belonging to other mussel species are also relevant (Sousa et al. 2008; Ilarri and Sousa 2012; Modesto et al. 2019).

Considering the high invasive potential of *C. fluminea* and its consequent economic and ecological impacts, it is important to understand the patterns of dispersion in non-native areas, accessing its updated distribution. This can help in predicting potential sites for new invasions and supporting preventive and controlling strategies, as well as identifying the main routes of introduction (Karatayev et al. 2007; Ilarri and Sousa 2012; Silveira et al. 2016). The objective of this study is to gather information on the current distribution of this species in Brazil and to record the occurrence of the species in the state of Piauí. Consequently, new elements to understand the process of introduction over time and the expansion of this mollusk in the country are provided.

Methods

Specimens were collected monthly between June 2017 and September 2019 in the urban perimeter of the Itaim River, in the city of Itainópolis, in southeastern Piauí. The 200 km-long Itaim River belongs to the Parnaíba river basin and covers the municipalities of Paulistana, Patos do Piauí, Itainópolis, Santa Cruz do Piauí, Paquetá, and Oeiras. The collection area is located in the semi-arid region of the country characterized by low rainfall, high evaporation rates, and high temperatures throughout the year (Ab'Sáber 2003; Silva et al. 2020).

The five sampling stations were located on the urban stretch of the river. The sampling was conducted using a sieve (21cm diameter) attached to a 1.5 m-long wooden rod. The sieve was dipped five times into the bottom of the river to take the sediment, and the specimens were then removed from the sieve using forceps. The separation and identification were preliminarily carried out considering the external and internal morphology of the shell, following Simone (2006), Mansur (2012), and Almeida et al. (2015). The *C. fluminea* individuals were preserved in 98% ethanol. Specimen lots were sent to the Museu de Zoologia da Universidade de São Paulo, São Paulo, Brazil (MZUSP) to confirm the identification. Field samplings were authorized by the Chico Mendes Institute for Biodiversity Conservation (ICMBio), through the Biodiversity Authorization and Information System (SISBIO), permit number 60423-5.

Information on the current distribution of *C. fluminea* in Brazil was obtained through extensive bibliographic search using the keywords *Corbicula fluminea* and Brazil, in the Google Scholar databases, National Center for Biotechnology Information (NCBI), Online Scientific Electronic Library (Scielo), Springer, and Science Direct. We selected scientific articles, theses, dissertations, and scientific meeting proceedings. Documents that did not mention vouchers deposited in scientific collections and/or how the specimens were identified were excluded from the research. The search also gathered data from specimens deposited in the Mollusca collection of the Museu de Zoologia da Universidade de São Paulo (MZUSP). The geographic coordinates available in the research sources were used to create a distribution map. When not included in the publications, the center coordinates of the municipality were adopted as the geographic position. This method of reviewing the distribution of species based on published data and scientific collections has already been used to gather information on the distribution of other mollusk species (Coelho et al. 2018; Santana et al. 2013; Miyahira et al. 2020).

Four categories were established to determine the types of environments in which the species was recorded in Brazil:

i) natural lotic environments: brook, creek, stream, river, and waterfall
ii) natural lentic environments: lake and pond
iii) anthropized environments: mining pit, dam, and reservoir
iv) unusual environments: one in which the species is not usually found, such as beaches and shale pits

Some records in natural environments occurred within areas of environmental protection, classified at the federal, state, and municipal levels.

Results

***Corbicula fluminea* (O.F. Müller, 1774)**

New record. BRAZIL • 3 living specimens; state of Piauí, Itainópolis, Itaim River; 07°26.54′S, 041°28.35′W; July 2019, 18 cm depth, sandy substrate; August 2019, 14 cm depth, sandy substrate and September 2019, 16 cm depth, sandy substrate; 1 specimen MZUSP 151417; Leal, M.F. (Fig. 1).

In total, 382 records were obtained in the bibliographic survey, with greatest number of records in the Southeast Region (n = 242, 63.4%), followed by the South (n = 57, 14.9%), Midwest (n = 44, 11.5%), North (n = 22, 5.8%) and Northeast regions (n = 17, 4.5%). There are records for *C. fluminea* in 179 municipalities belonging to 22 Brazilian states in addition to the Federal District (Fig. 2). This species occurs in all states of the South, Southeast, and Midwest regions. In the North Region, it occurs in four of the seven states and in the Northeast Region, in six of the nine states. Our new record of *C. fluminea* from Piauí is the first from this state and the Parnaíba river basin. With the addition of the Parnaíba river basin, *C. fluminea* is now known from every single river basin in the Brazilian territory (Fig. 2).

The highest portion of records occurred in natural lotic environments (n = 298, 78%), of which eight (2%) were within an area under the influence of a hydroelectric plant, followed by lentic natural environments (n = 70, 18.3%) and anthropized environments (n = 13, 3.4%) (Fig. 3). There was a single record (0.3%) from a beach (Fig. 3), which we classified as an unusual environment, probably a result of transportation. The examined literature indicated that 11 (3%) of the 368 records of *C. fluminea* in natural environments occurred inside environmental protection areas (Fig. 3).

Two (0.5%) of the 382 records of the species in Brazil did not present any information about the year of occurrence. The vast majority of records date from 2000, with special attention to 2019, which accumulated about a quarter of the records (n = 80) (Fig. 4). Malacological studies (n = 306, 80.1%) concentrate on the highest number of occurrences of *C. fluminea*. However, this species was also reported indirectly by research focusing on ecotoxicology (n = 47, 12.3%), aquatic invertebrates in general (n = 17, 4.5%), and ichthyology (n = 12, 3.1%).

Identification. The collected specimens of *C. fluminea* (Fig. 1) correspond to the descriptions given by Mansur and Pereira (2006) and Simone (2006); the triangular shell presents concentric striations on the external surface, generally a bright, yellowish-brown periostracum, an elevated and inflated umbo, and no pallial sinus. Variations in the internal shell color can occur (Foster et al. 2020) and are associated with environmental conditions (Ituarte 1994; Mansur 2012) and genetic variation (Qiu et al. 2001).



Figure 1. *Corbicula fluminea* specimen collected in the Itaim River, municipality of Itainópolis, Brazil. (MZUSP 151417, H 16 mm, L 19 mm). **A**. View of the outer surface, **B.** View of the inner surface. Scale bar = 10 mm.



Figure 2. Geographic distribution of *Corbicula fluminea* in Brazil, including the new record for the state of Piauí.



Figure 3. *Corbicula fluminea* records according to environment type in Brazil.

Discussion

The data presented in this work show that in the period of 40 years, *Corbicula fluminea* expanded its presence in the country, reaching 190 municipalities in all regions and all hydrographic basins, including the isolated Parnaíba river basin.

The high adaptive capacity of *C. fluminea* is proven by the diversity of environments in which it occurs (Sousa et al. 2008). High water temperature, shallow bed, and sandy substrate have been mentioned as ideal conditions to establish its populations (Zilli and Marchese 2011; Santana et al. 2013; Cao et al. 2017; Gama et al. 2017). These features are similar to the observed in the



Figure 4. *Corbicula fluminea* recorded in Brazil, between 1981 and 2019.

Itaim River and most rivers of the semi-arid region. The occurrence of *C. fluminea* in this type of environment threatens native species and aquatic biodiversity, considering that most tributaries are intermittent, with frequent decreases their volume and flow of water. Thus, the semi-arid region is favorable to a population increase of this exotic species.

Records of *C. fluminea* in lotic, lentic, anthropogenic, and unusual environments demonstrate the high invasiveness of this mollusk and its risk to environments and biological communities within them. As verified in our research, the greatest number of occurrences are in lotic environments in Brazilian territory. This is expected and deserves attention as running waters have a greater capacity to easily disperse these bivalves to new areas (Poleze and Callil 2015; Rosa and Dantas 2020). Similarly, the greater occurrence number in natural environments, whether lotic and lentic, indicate significant stress on native fauna, considering environmental changes caused by high population densities of *C. fluminea*, as discussed by Sousa et al. (2008), Silva and Stuff (2011), and Sazima and D'Angelo (2013). Thus, actions to control and mitigate this bioinvasion in Brazil are urgently needed.

The economic impact of *C. fluminea* has been calculated for several regions of the country (Mota 2006; Paschoal et al. 2013; Paschoal et al. 2015; Paschoal et al. 2018; Maroneze et al. 2011), but the impacts on the Northeast Region have not yet been assessed. As the intermittent water bodies in this region have great importance in supplying cities and rural communities, as well as providing for livestock, irrigation, and leisure, the presence of *C. fluminea* in high densities might compromise water quality and its use. The transport of water for human use, which due to its scarcity is common in the semi-arid Northeast Region, may favor the dispersion of this and other exotic aquatic species between unconnected watersheds.

The main routes that have led to the rapid expansion of *C. fluminea* in Brazil are still not well documented because aquarium practices, transport by birds, sand extraction, and pisciculture may also have contributed to this process (Gomes et al. 2016; Belz et al. 2012). Regarding the pisciculture, Coelho et al. (2018) have already noted it as an efficient agent for dispersal. The compiled results of our study show a large increase in the number of records of *C. fluminea* from the 2000s onwards. These years coincide with the growth pisciculture activities in Brazil according to Schulter and Vieira Filho (2017). Also of importance, although species in the family Cyrenidae do not have fish-parasitizing larvae like the Unionoida, they still can use fish as a means of passive transportation for upstream dispersion in rivers. Therefore, it is common to find live specimens in the intestines of catfish (Loricariidae) (Simone personal observation), which evacuate them alive.

The continuous expansion of the distribution of *C. fluminea* in Brazilian water bodies, calls attention to the need for studies on population dynamics, modeling of habitat adequacy, as well as genetic variability of populations. These researches must be considered for the development of strategies to avoid the advance of new environments as well as to mitigate possible damages to the native fauna (Pigneur et al. 2013; Silveira et al. 2016; Fernandes et al. 2020).

Acknowledgements

We are grateful to the Coordination of Superior Level Staff Improvement (CAPES) and to the National Council for Scientific and Technological Development (CNPq; 421582/2018–9) for providing financial support, and also to the Federal Institute of Piauí (PROAGRUPAR-INFRA 154/2019) for the logistic and financial support. We are also grateful to the anonymous reviewers and subject editor for the important contributions during the peer review process.

Authors contributions

MFL was responsible for the sample collection in the Itaim River, data analysis, and writing of the manuscript.

LRLS was responsible for the sample identification process and manuscript revision. ACFL coordinated data analysis and manuscript elaboration. ELS and TGP coordinated the research and revised the manuscript. All authors corrected, reviewed, and discussed the data.

References

Ab'Sáber AN (2003) Os domínios de natureza no Brasil: potencialidades paisagísticas. Ateliê Editorial, São Paulo, 159 pp.

Almeida ACC, Paz MCP, Paz RJ (2015) Occurrence of *Corbicula fluminea* (Müller, 1774) and *Corbicula largillierti* (Philippi, 1844) (Bivalve: Corbiculidae) in Municipality of Ingá (state of Paraíba, Northeast Brazil). Brazilian Journal of Biological Sciences 14 (2): 381–386. http://doi.org/10.21472/bjbs.020420

Belz CE, Darrigran G, Netto OSM, Boeger WA, Ribeiro Junior PJ (2012) Analysis of four dispersion vectors in inland waters: the case of the invading bivalves in South America. Journal of Shellfish Research 31 (3): 777–784. https://doi.org/10.2983/035.031.0322

Cao L, Damborenea C, Penchaszadeh PE, Darrigran G (2017) Gonadal cycle of *Corbicula fluminea* (Bivalvia: Corbiculidae) in Pampean streams (southern Neotropical Region). Plos ONE 12 (10): e0186850. https://doi.org/10.1371/journal.pone.0186850

Coelho PN, Fernandez MA, Cesar DAS, Ruocco AMC, Henry R (2018) Updated distribution and range expansion of the gastropod invader *Melanoides tuberculata* (Müller, 1774) in Brazilian waters. BioInvasions Records 7 (4): 405–409. https://doi.org/10.3391/bir.2018.7.4.08

Crespo D, Dolbeth M, Leston S, Sousa R, Pardal MA (2015) Distribution of *Corbicula fluminea* (Müller, 1774) in the invaded range: a geographic approach with notes on species traits variability. Biological Invasions 17 (1): 2087–2101. https://doi.org/10.1007/s10530-015-0862-y

Fernandes MR, Miyahira IC, Caetano CHS, Salgueiro F (2020) The spreading of the invasive bivalve *Mytilopsis leucophaeata* (Dreissenidae) into estuaries of Rio de Janeiro, Brazil. Biological Sciences 92 (Supplement 2): 1–12. https://doi.org/10.1590/0001-3765202020190045

Foster A, Fuller P, Benson A, Constant S, Raikow D, Larson J, Fusaro A (2020) *Corbicula fluminea*. USGS nonindigenous aquatic species database, Gainesville, USA. https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=92. Accessed on: 2020-11-11.

Freire CG, Marafon AT (2018) Invasive mollusc species in Brazilian aquatic ecosystems and their impact on the environment. InterfacEHS - Saúde, Meio Ambiente e Sustentabilidade 13 (1): 2–16.

Gama M, Crespo D, Dolbeth M, Anastácio PM (2017) Ensemble forecasting of *Corbicula fluminea* worldwide distribution: Projections of the impact of climate change. Aquatic Conservation: Marine and Freshwater Ecosystems 27 (3): 675–684. https://doi.org/10.1002/aqc.2767

Gama M, Crespo D, Dolbeth M, Anastácio P (2016) Predicting global habitat suitability for *Corbicula fluminea* using species distribution models: The importance of different environmental dataset. Ecological Modelling 319: 163–169. https://doi.org/10.1016/j.ecolmodel.2015.06.001

Gomes C, Sousa R, Mendes T, Borges R, Vilares P, Vasconcelos V, Guilhermino L, Antunes A (2016) Low genetic diversity and high invasion success of *Corbicula fluminea* (Bivalvia, Corbiculidae) (Müller, 1774) in Portugal. Plos ONE 11 (7): e0158108. https://doi.org/10.1371/journal.pone.0158108

Ilarri MI, Sousa R (2012) *Corbicula fluminea* Müller (Asian clam). In: Francis RA (Ed.) A handbook of global freshwater invasive species. Earthscan, London, UK, 173–183 pp.

Ituarte CF (1994) *Corbicula* and *Neocorbicula* (Bivalvia: Corbiculidae) in the Paraná, Uruguay, and Río de La Plata basins. The Nautilus 107 (4): 129–136.

Karatayev A, Padilla DK, Minchin D, Boltovskoy D, Burlakova LE (2007) Changes in global economies and trade: the potential spread of exotic freshwater bivalves. Biological Invasions 9 (2): 161–180. https://doi.org/10.1007/s10530-006-9013-9

Lockwood JL, Hoopes M, Marchetti MP (2007) Invasion ecology. Blackwell Publishing, Singapore, 304 pp.

Magalhães PS, Simões NR, Sonoda SL (2016) Limnologia de rios intermitentes: a bacia hidrográfica do rio Jequiezinho como estudo de caso. In: Moraes MEB, Lorandi R (Eds.) Métodos e técnicas de pesquisa em bacias hidrográficas. Editora Editus, Ilhéus, Brazil, 163–181 pp.

Mansur MCD (2012) Bivalves invasores límnicos: morfologia comparada de *Limnoperna fortunei* e espécies de *Corbicula* spp. In: Mansur MCD, Santos CP, Pereira D, Paz ICP, Zurita MLL, Rodriguez MTR, Nehrke MV, Bergonci PEA (Eds.) Moluscos límnicos invasores no Brasil: biologia, prevenção e controle. Redes Editora, Porto Alegre, Brazil, 61–74 pp.

Mansur MCD, Pereira D (2006) Bivalves límnicos da bacia do Rio dos Sinos, Rio Grande do Sul, Brasil (Bivalvia, Unionoida, Veneroida e Mytiloida). Revista Brasileira de Zoologia 23 (4): 1123–1147. https://doi.org/10.1590/S0101-81752006000400021

Maroneze DM, Coscarelli D, Vidigal THDA, Callisto M (2011) First record of *Corbicula fluminea* (Müller, 1774) in the drainage basin of the Araguari River, Minas Gerais, Brazil. Brazilian Journal of Biology 71 (1): 221–222. https://doi.org/10.1590/S1519-69842011000100033

Martins DS, Veitenheimer-Mendes IL, Faccioni-Heuser MC (2006) Aspectos morfológicos e de incubação em três espécies de *Corbicula* Mühlfeld, no Lago Guaíba, Rio Grande do Sul, Brasil (Bivalvia, Corbiculidae). Biota Neotropica 6 (2): 1–11.

Meyer AAN, Oliveira E, Silva IS, Prestes JG (2017) Evaluation of density of freshwater bivalves from the Environmental Protection Area (APA) of Verde River, Campo Magro, Paraná, Brazil. Pan-American Journal of Aquatic Sciences 12 (1): 21–30.

Miyahira IC, Pereira LS, Santos LN (2020) Non-native freshwater molluscs in the Neotropics: what can be learned from Brazilian reservoirs? Aquatic Invasions 15 (3): 455–472. https://doi.org/10.3391/ai.2020.15.3.06

Modesto V, Castro P, Lopes-Lima M, Antunes C, Ilarri M, Sousa R (2019) Potential impacts of the invasive species *Corbicula fluminea* on the survival of glochidia. Science of the Total Environment 673: 157–164.

Mota RH (2006) Análise da influência da geoquímica do ambiente e das características do substrato na estruturação da população de *Corbicula fluminea*, Muller 1974 (mollusca, bivalvia) no reservatório da usina hidrelétrica de Molta Grande–MG/SP. Dissertation, Fundação Universidade Federal de Ouro Preto, Ouro Preto, Brazil, 94 pp.

Paschoal LRP, Andrade DA, Cavallari DC (2013) First record of *Aylacostoma francana* (Ihering, 1909) (Gastropoda, Thiaridae) in Minas Gerais state, Brazil. Revista Biotemas 26 (2): 277–281.

Paschoal LRP, Andrade DP, Darrigran G (2015) How the fluctuations of water levels affect populations of invasive bivalve *Corbicula fluminea* (Müller, 1774) in a Neotropical reservoir? Brazilian Journal of Biology 75 (1): 135–143. https://doi.org/10.1590/1519-6984.09113

Paschoal LRP, Lacerda LEM, Miyahira IC (2018) Taking a ride! Phoretic association of the freshwater limpet *Gundlachia radiata* (Planorbidae: Ancylinae) and the apple snail *Pomacea lineata* (Caenogastropoda: Ampullariidae), with the description of a new record of this ancylinid in southeastern Brazil. Pan-American Journal of Aquatic Sciences 13 (4): 306–310.

Pfeifer NTS, Pitoni VLL (2003) Análise qualitativa estacional da fauna de moluscos límnicos no Delta do Jacuí, Rio Grande do Sul, Brasil. Biociências 11 (2): 145–158.

Pigneur LM, Falisse E, Roland K, Everbecq E, Deli ege JF, Smitz JS, Doninck KV, Descy JP (2013) Impact of invasive Asian clams, *Corbicula* spp., on a large river ecosystem. Freshwater Biology 59 (3): 1–11.

Pimpão DM, Martins DS (2009) Moluscos de água doce do Tupé, Manaus, AM, Brasil. In Silva ENS, Scudeller VV (Eds) Biotupé: Meio físico, diversidade biológica e sociocultural do Baixo Rio Negro, Amazônia Central volume 2. UEA Edições, Manaus, Brazil, 63–70.

Poleze M, Callil CT (2015) Bivalvia, Cyrenidae, *Corbicula fluminea* (Müller, 1774): new record, density, and population structure in the Teles Pires River, northern Mato Grosso, Brazil. Check List 11 (4): 1720. https://doi.org/10.15560/11.4.1720

Qiu A, Shi A, Komaru A (2001) Yellow and brown shell color morphs of *Corbicula fluminea* (Bivalvia: Corbiculidae) from Sichuan Province, China, are triploids and tetraploids. Journal of Shellfish Research 20 (1): 323–328.

Ragonha FH, Pinha GD, Petsch DK, Mansur MCD, Takeda AM (2014) First records of freshwater bivalves of Ilha Grande National Park, Paraná, Brazil. Iheringia, Série Zoologia 104 (1): 14–20. http://doi.org/10.1590/1678-4766201410411420

Rosa LC, Dantas JO (2020) First record of the Asian clam *Corbicula fluminea* (Müller, 1774) (Bivalvia: Cyrenidae) at Poxim-Açu River, northeastern Brazil. Acta Limnologica Brasiliensia 32: e22. https://doi.org/10.1590/s2179-975x8019

Santana DO, Silva MJM, Bocchiglieri A, Pantaleão SM, Faria RG, Souza BB, Rocha SM, Lima LFO (2013) Mollusca, Bivalvia, Corbiculidae, *Corbicula fluminea* (Müller, 1774): First record for the Caatinga biome, northeastern Brazil. Check List 9 (5): 1072–1074. http://doi.org/10.15560/9.5.1072

Santos SB, Thiengo SC, Fernandez MA, Miyahira IC, Gonçalves ICB, Ximenes RF, Mansur MCD, Pereira D (2012) Espécies de moluscos límnicos invasores no Brasil. In: Mansur MCD, Santos CP, Pereira D, Paz ICP, Zurita MLL, Rodriguez MTR, Nehrke MV, Bergonci PEA (Eds) Moluscos límnicos invasores no Brasil: biologia, prevenção e controle. Redes editora, Porto Alegre, Brazil, 61–74.

Sazima I, D'Angelo GB (2013) The Asian invasive freshwater clam *Corbicula fluminea* as prey of two native waterbirds in South-Eastern Brazil. Folia Malacologica 21 (4): 293–295. https://doi.org/10.12657/folmal.021.031

Schulter EP, Vieira Filho JER (2017) Evolução da piscicultura no Brasil: diagnóstico e desenvolvimento da cadeia produtiva de tilápia. Instituto de pesquisa econômica aplicada, Brasília, 42 pp.

Silva EL, Rocha AJ, Leal MF, Santos O, Sousa JH, Silva ARV, Dantas KKS, Matos EM, Castro ES, Pacheco ACL, Pinheiro TG (2020) Freshwater mollusks from three reservoirs of Piauí, northeastern Brazil. Biota Neotropica 20 (1): 1–8. https://doi.org/10.1590/1676-0611-bn-2019-0868

Silva LG, Stuff T (2011) Distribuição e abundância dos bivalves *Corbicula fluminea* (Mueller, 1774) e *Neocorbicula limosa* (Maton, 1811) no lago Guaíba, Rio Grande do Sul, Brasil. Fepam em Revista 5 (1): 13–17.

Silveira TCL, Gama AMS, Alves TP, Fontoura NF (2016) Modeling habitat suitability of the invasive clam *Corbicula fluminea* in a Neotropical shallow lagoon, southern Brazil. Brazilian Journal of Biology 76 (3): 718–725.

Simberloff D (2009) The Role of propagule pressure in biological invasions. Annual Review of Ecology, Evolution, and Systematics 40 (1): 81–102. https://doi.org/10.1146/annurev.ecolsys.110308.120304

Simone LRL (2006) Land and freshwater Molluscs of Brazil. Fapesp, São Paulo, Brazil, 390 pp.

Sousa R, Antunes C, Guilhermino L (2008) Ecology of the invasive Asian clam *Corbicula fluminea* (Müller, 1774) in aquatic ecosystems: an overview. Annales de Limnologie - International Journal of Limnology 44 (2): 85–94. https://doi.org/10.1051/limn:2008017

Veitenheimer-Mendes IL (1981) *Corbicula manilensis* (Philippi, 1844) molusco asiático, na bacia do Jacuí e do Guaíba, Rio Grande do Sul, Brasil (Bivalvia, Corbiculidae). Iheringia, Série Zoológica 60 (1): 63–74.

Zilli FL, Marchese MR (2011) Patterns in macroinvertebrate assemblages at different spatial scales. Implications of hydrological connectivity in a large floodplain river. Hydrobiologia 663 (1): 245–257. https://doi.org/10.1007/s10750-010-0576-1